

PRE-OPERATIVE INSTRUCTIONS

RIGHT

LEFT



Acceptable Substitutions:
Gatifloxacin, Vigamox or Moxifloxacin






Acceptable Substitutions:
Pred Forte






Acceptable Substitutions:
Bromfenac, Ilevro or Bromsite

1 DAY PRIOR TO SURGERY

 BESIVANCE	<input type="checkbox"/> Breakfast <input type="checkbox"/> Lunch <input type="checkbox"/> Bedtime
 DUREZOL	<input type="checkbox"/> Breakfast <input type="checkbox"/> Lunch <input type="checkbox"/> Bedtime
 PROLENSA	<input type="checkbox"/> Bedtime

DAY OF SURGERY

 BESIVANCE	<input type="checkbox"/> Before going to hospital <input type="checkbox"/> 2 hours after returning home from hospital <input type="checkbox"/> Bedtime
 DUREZOL	<input type="checkbox"/> Before going to hospital <input type="checkbox"/> 2 hours after returning home from hospital <input type="checkbox"/> Bedtime
 PROLENSA	<input type="checkbox"/> Bedtime

*Mark out a circle for each drop you take

*The order you put drops in your eye does not matter, but be sure you space each medicine 5 minutes apart

*Shake bottle prior to use

Medication Instruction Sheet



Georgia Eye
ASSOCIATES

**Donald E. Poland, MD • Kris F. Gillian, MD • Dickie McMullan, MD • Benjamin Strauss, MD
Frank L. Winski, OD • Brigette Rabitsch, OD • Brian A. Kahn, OD**

Patient Name: _____

Surgery Date: _____

Operation Eye: Right / Left Eye

Special Instructions:

- 1.) Do not eat or drink anything (including coffee) after midnight, on the day before your surgery.
- 2.) Any oral medicines that you normally take in the morning may be taken with a very small sip of water, on the morning of your surgery. All patients with diabetes should contact their internist on how to manage their insulin on the day of surgery.
- 3.) Stop all aspirin (or aspirin containing products) 2 weeks prior to your surgery. All blood thinners such as: coumadin, plavix, celebrex should be stopped three days prior to your surgery. You may resume blood thinners immediately after surgery (same day).
- 4.) Do not wear makeup the day of your surgery.
- 5.) You must have a driver to drive you home after surgery.
- 6.) Keep tip of eye dropper clean and do not touch tip to the eye or any other object.
- 7.) Wait 5 minutes between each different eye medication.
- 8.) Each medication has refills if needed.

Lawrenceville
771 Old Norcross Road
Suite 150
Lawrenceville, GA 30045
770.995.5408
Faz: 770.513.2042

Atlanta
3120 Maple Drive
Atlanta, GA 30305
404.233.3267
Fax: 404.233.4399

Tucker
2368 Main Street
Suite B
Tucker, GA 30084
770.938.0020
Fax: 770.939.1256